


NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




Breakpoint = 64

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates


Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)



NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates


Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)



NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




Breakpoint = 4

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




*Breakpoint Not Established

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates


Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)



NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)




↓ Breakpoint

NARMS – EB 1999

Veterinary Isolates

Fig. 22. Minimum Inhibitory Concentrations by Antimicrobial Agent Major Serotypes from Cattle (Diagnostic)



↓ Breakpoint